CLAIMS

1. A compound of formula

$$X-Y$$
 $(CH_2)_q$ R^4 R^6 R^7 $(R^3)_t$ $(R^2)_n$ $(R^2)_n$ $(R^2)_n$ $(R^3)_t$ $(R^3)_t$

wherein

m is 0, 1, 2, 3 or 4;

each R^1 independently represents halogen, cyano, hydroxyl, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 alkoxy or sulphonamido;

X represents a bond, -CH₂- or -O-, Y represents a bond, -CH₂- or -O-, and Z represents a bond, -O-, -NH- or -CH₂-, provided that only one of X, Y and Z can represent a bond at any one time and provided that X and Y do not both simultaneously represent -O-;

n is 0, 1 or 2;

each R² independently represents halogen, C₁-C₆ alkyl or C₁-C₆ haloalkyl;

q is 0 or 1;

15

20

t is 0, 1, 2, 3, 4 or 5;

each R³ independently represents halogen, cyano, nitro, hydroxyl, -C(O)H, -NR⁹R¹⁰, -CH₂C(O)NR¹¹R¹², -CH₂NHC(O)R¹³, -NHSO₂R¹⁴, -SO₂NR¹⁵R¹⁶, -CH₂-R¹⁷, C₁-C₆ alkylcarbonyl, phenylcarbonyl, C₃-C₆ cycloalkyl, or a group selected from C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkoxy, phenyl and a saturated or unsaturated 5- to 10-membered heterocyclic ring system comprising at least one ring heteroatom selected from nitrogen, oxygen and sulphur, each group being optionally substituted with at least one substituent selected from halogen, cyano, hydroxyl, carboxyl, C₁-C₆ alkyl, C₁-C₆ alkoxy and C₁-C₆ alkoxycarbonyl;

15

 R^4, R^5, R^6, R^7 and R^8 each independently represent hydrogen, halogen, C_1 - C_6 alkyl or C_1 - C_6 haloalkyl;

R⁹ and R¹⁰ each independently represent hydrogen or C₁-C₆ alkyl;

R¹¹ and R¹² each independently represent hydrogen or C₁-C₆ alkyl, or R¹¹ and R¹²

together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocyclic ring which may be optionally substituted with at least one substituent selected from hydroxyl;

R¹³ and R¹⁴ each independently represent hydrogen or C₁-C₆ alkyl; and

R¹⁵ and R¹⁶ each independently represent hydrogen or C₁-C₆ alkyl, or R¹⁵ and R¹⁶

together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocyclic ring which may be optionally substituted with at least one substituent selected from hydroxyl;

R¹⁷ is a 5 to 7 membered saturated heterocyclic ring containing at least one nitrogen atom, which ring may be optionally substituted with one or more oxo groups; or a pharmaceutically acceptable salt or solvate thereof.

2. A compound according to claim 1, wherein X and Y have the meanings shown in the following table:

| X | Y |
|-----------------|-----------------|
| bond | 0 |
| 0 | bond |
| CH ₂ | bond |
| bond | CH ₂ |

- 3. A compound according to claim 1 or claim 2, wherein Z represents -O- or -CH₂-.
 - 4. A compound according to any one of claims 1 to 3, wherein q is 1.

WO 2005/049620 PCT/SE2004/001658

- 5. A compound according to any one of claims 1 to 4, wherein m is 1 and R¹ represents halogen.
- 6. A compound according to any one of claims 1 to 5, wherein each R³ independently represents halogen, cyano, nitro, hydroxyl, -C(O)H, -NR⁹R¹⁰, -CH₂C(O)NR¹¹R¹², -CH₂NHC(O)R¹³, -NHSO₂R¹⁴, -SO₂NR¹⁵R¹⁶, -CH₂-R¹⁷, C₁-C₄ alkylcarbonyl, phenylcarbonyl, C₅-C₆ cycloalkyl or a group selected from C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ alkoxy, phenyl and a saturated or unsaturated 5- to 6-membered heterocyclic ring system comprising one, two, three or four ring heteroatoms independently selected from nitrogen, oxygen and sulphur, each group being optionally substituted with one, two, three or four substituents independently selected from halogen, cyano, hydroxyl, carboxyl, C₁-C₄ alkyl, C₁-C₄ alkoxy and C₁-C₄ alkoxycarbonyl.
- 7. A compound according to claim 6, wherein the saturated or unsaturated 5- to 6-membered heterocyclic ring system is isoxazolyl, pyrrolyl, morpholinyl, piperidinyl or oxadiazolyl.
 - 8. A compound according to claim 1 selected from:

10

20

25

30

(2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(2-methoxyphenoxy)propan-2-ol hydrochloride,

2-{[(2S)-3-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy}phenol,

(2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-[2-(2-hydroxyethoxy)phenoxy]propan-2-ol hydrochloride,

2-(2-{[(2S)-3-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy}phenyl)-N-methylacetamide trifluoroacetate (salt),

(3S)-1-[(2-{[(2S)-3-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy}phenyl)acetyl]pyrrolidin-3-ol,

N-(2-{[(2S)-3-(5-Chloro-1'H.3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl-2-hydroxypropyl]oxy}benzyl)acetamide,

20

- 2-(2-{[(2S)-3-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy}-4-methoxyphenyl)-N-methylacetamide,
- 2-(2-{[(2S)-3-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'yl)-2-hydroxypropyl]oxy}-4-hydroxyphenyl)-N-methylacetamide trifluoroacetate (salt),
- 2-(4-{[(2S)-3-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy}-2-methoxyphenyl)-N-methylacetamide,
- (2S)-1-(2-Amino-5-methoxyphenoxy)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)propan-2-ol bis(trifluoroacetate),
- $N-(2-\{[(2S)-3-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy\}-4-hydroxyphenyl) methanesulfonamide trifluoroacetate,$
- $N-(2-\{[(2S)-3-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy\}-4-methoxyphenyl) methanesulfonamide trifluoroacetate,$
- (2S)-1-(4-Bromo-2-fluorophenoxy)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)propan-2-ol,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(3-ethynylphenoxy)propan-2-ol,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(2,4-dichloro-3,5-dimethylphenoxy)propan-2-ol,
- (2S)-1-(4-Chloro-2-isoxazol-5-ylphenoxy)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)propan-2-ol,
 - (4-{[(2S)-3-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy}phenyl)(phenyl)methanone,
 - (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(2,3,4,6-tetrachlorophenoxy)propan-2-ol,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(2-cyclohexyl-5-methylphenoxy)propan-2-ol,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]=1'-yl)-3-phenoxypropan-2-ol,
- (2S)-1-(2-Bromophenoxy)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)propan-2-ol,

15

20

- 2-{[(2S)-3-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy}benzaldehyde,
- 5-tert-Butyl-2-{[(2S)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy}benzaldehyde,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(1,1':3',1"-terphenyl-2'-yloxy)propan-2-ol,
- 1-(2-{[(2S)-3-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy}-5-methoxyphenyl)ethanone,
- 1-(5-Bromo-2-{[(2S)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy}phenyl)ethanone,
 - (2S)-1-(4-Chloro-2-isopropyl-5-methylphenoxy)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)propan-2-ol,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(2,3-dimethyl-4-nitrophenoxy)propan-2-ol,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(2,4-dichlorophenoxy)propan-2-ol,
- Ethyl (2E)-3-(4-{[(2S)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy}-3-methoxyphenyl)acrylate,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(2-methyl-3-nitrophenoxy)propan-2-ol,
- 5-Chloro-2-{[(2S)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy}benzaldehyde,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(2-fluorophenoxy)propan-2-ol,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(3-fluorophenoxy)propan-2-ol,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(4-fluorophenoxy)propan-2-ol,
- (2S)-1-(2-Chlorophenoxy)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)propan-2-ol,

15

20

- (2S)-1-(3-Chlorophenoxy)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)propan-2-ol,
- (2S)-1-(4-Chlorophenoxy)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)propan-2-ol,
- (2S)-1-(3-Bromophenoxy)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)propan-2-ol,
 - (2S)-1-(4-Bromophenoxy)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)propan-2-ol,
 - (2S)-1-(2-tert-Butyl-5-methylphenoxy)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)propan-2-ol,
 - (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-[2-(trifluoromethyl)phenoxy]propan-2-ol,
 - 1-(2-{[(2S)-3-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy}-4,5-dimethoxyphenyl)ethanone,
 - (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]propan-2-ol,
 - (2S)-1-(4-Chloro-3-ethylphenoxy)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)propan-2-ol,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-[3-(2,5-dimethyl-1H-pyrrol-1-yl)phenoxy]propan-2-ol,
 - (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-[2-(hydroxymethyl)phenoxy]propan-2-ol,
 - (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-[2-(2-hydroxyethyl)phenoxy]propan-2-ol,
 - 3-{[(2S)-3-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy}benzonitrile,
 - 2-{[(2S)-3-(5-Chlörö-1'H,3H-spiró[1-benzofuran-2,4'-piperidin]-1'-yl)-2- hydroxypropyl]oxy}benzonitrile,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(2-morpholin-4-yl)-30 ylphenoxy)propan-2-ol,

- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(2,3-difluoro-6-nitrophenoxy)propan-2-ol,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-y1)-3-(2,3,6-trichlorophenoxy)propan-2-ol,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(4-fluoro-2-methoxyphenoxy)propan-2-ol,
 - 5-Chloro-2-{[(2S)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy}-3-methylbenzaldehyde,
 - (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-[4-(4-methylpiperidin-1-yl)-2-nitrophenoxy]propan-2-ol,
 - (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(2,4-dichloro-3,5-dimethyl-6-nitrophenoxy)propan-2-ol,
 - 1-(3,5-Dichloro-2-{[(2S)-3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy}phenyl)propan-1-one,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(4-ethylphenoxy)propan-2-ol,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(2-ethylphenoxy)propan-2-ol,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(3-ethylphenoxy)propan-2-ol,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-(3-morpholin-4-ylphenoxy)propan-2-ol,
- (2S)-1-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-[2-(5-methyl-1,3,4-oxadiazol-2-yl)phenoxy]propan-2-ol,
- 4-{[(2S)-3-(5-Chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2-hydroxypropyl]oxy}benzonitrile,
 - (2S)-1-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-3-[2-(pyrrolidin-1-ylsulfonyl)phenoxy]propan-2-ol.
- 1-(2-{[(2S)-3-(5-chloro-1'*H*, 3*H*-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)2hydroxypropoxy]benzyl}imidazoline-2,4-dione,

(IV)

- (2S)-{2-chloro-5-[3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2hydroxypropoxy]phenoxy}acetic acid,
- (2S)-{2,4-dichloro-5-[3-(5-chloro-1'H,3H-spiro[1-benzofuran-2,4'-piperidin]-1'-yl)-2hydroxypropoxy]phenoxy}acetic acid,
- and pharmaceutically acceptable salts and solvates of any one thereof.
 - A process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as defined in claim 1 which comprises,

(a) reacting a compound of formula

10

15

wherein m, R¹, n, R², q, X, Y and Z are as defined in formula (I), with a compound of formula

wherein $t, R^3, R^4, R^5, R^6, R^7$ and R^8 are as defined in formula (I); or

(b) reacting a compound of formula

$$(R^{1})_{m}$$

$$X-Y$$

$$(CH_{2})_{q}$$

$$R^{5}$$

$$R^{5}$$

$$R^{8}$$

$$R^{7}$$

$$(R^{2})_{n}$$

wherein m, R^1 , n, R^2 , q, X, Y, Z, R^4 , R^5 , R^6 , R^7 and R^8 are as defined in formula (I), with a compound of formula

10

15

HO
$$(R^3)_t$$
 (V)

wherein t and R³ are as defined in formula (I), in the presence of a suitable base; or

(c) when t is at least one and a group R³ represents -NHSO₂R¹⁴, reacting a compound of formula

$$(R^{1})_{m}$$
 $X-Y$
 $(CH_{2})_{q}$
 R^{5}
 R^{5}
 R^{7}
 $(R^{3})_{l'}$
 $(R^{2})_{n}$
 $(R^{2})_{n}$

wherein t' is 0, 1, 2, 3 or 4, R^3 ' is as defined for R^3 in formula (I) other than -NHSO₂ R^{14} and m, R^1 , n, R^2 , q, X, Y, Z, R^4 , R^5 , R^6 , R^7 and R^8 are as defined in formula (I), with a compound of formula

wherein L¹ represents a leaving group and R¹⁴ is as defined in formula (I), in the presence of a suitable base;

(d) where t is at least 1 and a group R³ represents -CH₂-R17, where R17 is a 5 to 7-membered saturated heterocyclic ring containing 2 nitrogen atoms and which ring is substituted by two oxo groups, reacting a compound of formula

10

15

$$(R^{1})_{m}$$

$$(R^{2})_{n}$$

$$(R^{2})_{n}$$

$$(VIII)$$

wherein t' is 0, 1, 2, 3 or 4, R³ is as defined for R³ in formula (I) other than -CH2-R17, and m, R¹, n, R², q, X, Y, Z, R⁴, R⁵, R⁶, R⁷ and R⁸ are as defined in formula (I), with an alkyl glycinate in the presence of a reducing agent, and subsequently with metal isocyanate;

and optionally after (a), (b) or (c) forming a pharmaceutically acceptable salt or solvate.

- 10. A pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1 to 8 in association with a pharmaceutically acceptable adjuvant, diluent or carrier.
- 11. A process for the preparation of a pharmaceutical composition as claimed in claim 10 which comprises mixing a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1 to 8 with a pharmaceutically acceptable adjuvant, diluent or carrier.
- 12. A compound of formula (I) or a pharmaceutically-acceptable salt or solvate thereof as claimed in any one of claims 1 to 8 for use in therapy.
- 13. Use of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1 to 8 in the manufacture of a medicament for the treatment of human diseases or conditions in which modulation of chemokine receptor activity is beneficial.

- 14. Use of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1 to 8 in the manufacture of a medicament for use in treating rheumatoid arthritis.
- 15. Use of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1 to 8 in the manufacture of a medicament for use in treating chronic obstructive pulmonary disease.
- 16. Use of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1 to 8 in the manufacture of a medicament for use in treating asthma.
 - 17. Use of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1 to 8 in the manufacture of a medicament for use in treating multiple sclerosis.
 - 18. A method of treating an inflammatory disease which comprises administering to a patient in need thereof a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1 to 8.
 - 19. A method of treating an airways disease which comprises administering to a patient in need thereof a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1 to 8.

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

| BLACK BORDERS . |
|---|
| IMAGE CUT OFF AT TOP, BOTTOM OR SIDES |
| FADED TEXT OR DRAWING |
| BLURRED OR ILLEGIBLE TEXT OR DRAWING |
| SKEWED/SLANTED IMAGES |
| ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS |
| GÉAY SCALE DOCUMENTS |
| LINES OR MARKS ON ORIGINAL DOCUMENT |
| REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY |
| □ OTHER: |

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.